

Original article

X-ray evaluation at 5–6 years of Straumann implants (part 2)

Hamasat Gheddaf-Dam¹, Semaan Abi Najm², Stella Lysitsa³, Mark Bischof², Rabah Nedir²

¹ Department of Prosthodontics and Operative Dentistry, Tufts School of Dental Medicine Boston, USA

² Division of Stomatology and Oral Surgery, Dental School, Geneva, Switzerland

³ Department of Oral Surgery, Aristotle University, Thessaloniki, Greece

Discussion

Long term preservation of crestal bone height around osseointegrated implants is often used as a measure of primary success [5, 12, 22]. Prospective long-term studies exhibited survival and success rates largely exceeding 95% after 5 and 10 years of follow-up for the Straumann® implant system [3, 5, 18, 22, 45, 57]. A mean crestal bone loss ≤ 1.5 mm during the first year and ≤ 0.2 mm per year thereafter is proposed as one of the major success criteria. If we apply these strict success criteria then the CBL in 5 years should not exceed 2.3 mm [$1.5 + (0.2 \times 4)$].

In the current study, 8.5% of the implants exhibited “supra-boundary bone”. In addition to 84.5% of the implants showing bone loss within the physiological range (0–3 mm), giving an overall successful pool of implants up to 93%. This represents a high success rate considering the private practice setting and the absence of exclusion criteria in the initial enrolment of the patients.

Patients with implants exhibiting a bone loss of 2–3 mm (7.8%) would require careful monitoring, with closer hygiene recalls and increased education in regard to patient awareness for dental hygiene and maintenance.

Bone loss greater than 3 mm was observed in 7% of the included implants. At the 5–6 year control, they were still well integrated in the jaw bone and the subjects did not manifest any symptoms that previously identified them as unsuccessful [7]. Moreover, the status and prognosis of such implants have to be carefully interpreted because other factors, mainly clinical parameters such as bleeding on probing and pocket depths, were not available, in contrary to previous studies [5, 12, 31, 38, 39, 45–47, 51]. Considering such arguments, and although these implants were put by the study group in the unsuccessful implants category, one could argue the contrary. This group deserves careful monitoring, with closer hygiene recalls, more follow-up radiographs, and extended patient awareness to dental hygiene.

For the Straumann® implants, the distance from the implant shoulder to the first bone–implant contact was called

DIB (distance implant–bone) and was used in previous studies [16, 21]. These studies followed the changes in peri-implant bone levels over time by taking measurement between two time points. A baseline and a postoperative radiograph were usually taken to identify initial and final bone levels, and therefore to calculate the difference: Δ DIB.

In the present study, an original method was used to calculate the bone change: the interface of the smooth-roughened surface (identified as the R interface) was considered as the baseline level. It was assumed to be the level up to which bone loss was considered as physiological, i.e. not affected by external factors. Bone loss occurring further from this point, in an apical implant direction, was thus identified as crestal bone loss (CBL). Bone localized coronally to this interface was defined as “supra-boundary bone”. When the bone level was stabilized at the interface, it was then considered that no bone change occurred (CBL and “supra-boundary bone” were equal to 0 mm).

From that point, it was interesting to identify factors that might enhance bone loss or favor bone maintenance when single-staged Straumann® implants with treated TPS/SLA surfaces were used. It could be argued that the mean CBL was even lower than the 1.2 mm obtained in this study, as this value was compensated with implants that had what was called “supra-boundary bone”. The present study did not quantify this “supra-boundary bone”, as was described in a recently published study [51].

A paralleling radiographical technique may sometimes be difficult to perform because of the implant inclination and patient anatomy. For example, in the case of an extremely resorbed mandible, the intra-oral placement of the film was impossible because of the interference of the mouth floor [33]. In the maxilla, where the palate is the most inclined, it was difficult to position the film without bending [34]. This explained the large number of un-interpretable radiographs in these two regions.

When observing the comparative tables for the statistically significant results, namely surface texture, smoking status, anterior/posterior location and VBL, it is worth noticing

that they strongly influenced both extremes of bone change groups, i.e. bone loss higher than 3 mm and bone occurring above the rough-smooth surface ("supra-boundary bone"). For example, a TPS-surfaced implant, an implant placed in a smoker, an implant localized in an anterior arch, and an implant with VBL < 1 mm showed higher differences than their counterparts in zones of bone loss > 3 mm (higher bone loss) or in zones of "supra-boundary bone" (least "supra-boundary bone"). Their influence became low in zones of physiological bone loss (0–3 mm).

Statistically significant factors affecting bone loss

Surface texture effect

In the past 15 years, the topography of titanium surfaces has been investigated for dental implant applications [7, 16, 21, 37, 56]. The main goal of these experimental studies was to determine whether bone apposition could be enhanced by new microrough titanium surfaces as compared with the original implant surfaces utilized in implant dentistry, such as machined or titanium-plasma-sprayed (TPS) surfaces. Various techniques have been used to produce microrough titanium surfaces, including sandblasting, acid etching, or combinations of those, to modify surface topography. SLA surfaces were shown to have greater success and survival rates than TPS surfaces in several animal (miniature pigs) studies and human studies [70–73].

Among these new surfaces, the sandblasted and acid-etched (SLA) surface demonstrated enhanced bone apposition in histomorphometric studies and higher removal torque values in biomechanical testing [71–74]. Based on these experimental results, clinical studies were initiated to load SLA implants after a reduced healing period of only 6 weeks. In the study of Cochran *et al.* [71] comparing TPS with SLA surfaces in the canine mandible, linear measurements on standardized radiographs from the implant shoulder to first bone-to-implant contact (DIB) were done and bone density was evaluated by computer-assisted densitometric image analysis (CADIA). DIB measurements indicated that SLA implants significantly showed less bone height loss (0.52 mm) than TPS implants (0.69 mm). Histometric findings by the same groups later confirmed these results. The SLA implants exhibited significantly higher percentage of bone-to-implant contact than did the TPS implants [72]. The clinical examination up to 3 years demonstrated favorable results, with success rates around 99% [71]. A more recent study revealed very high success rates for SLA surfaced implants loaded at 6 weeks and placed in the posterior maxilla [74].

To date, this study is the first to have equally large numbers of each surface. The present study also clearly yielded significant higher bone loss on the TPS implants than on the SLA surface implants. Moreover, the TPS group had a greater proportion of implants having bone loss > 3 mm. The TPS surface became more significant in terms of degrees of bone

loss when combined with other factors such as smoking, anterior location and VBL prior to implant placement smaller than 1 mm. The proportion of bone above the rough-machined interface (named "supra-boundary bone" in the current study) was also significantly higher on the SLA surface. Therefore, it may be prudent to establish a more intense oral hygiene follow-up for patients with TPS-surfaced implants, especially if other aggravating factors, such as tobacco use, are present.

Of note, however, is that TPS surfaced implants still osseointegrated well and have contributed to the high survival rate of the study. Hence, although they have greater crestal bone loss (lower success rates), TPS surfaced implants still represent a valid treatment option for implants in function.

Tobacco effect

The effect of tobacco on dental implants is well documented in the literature and many authors have shown that heavy smokers show a greater degree of crestal bone loss when compared to non-smokers [49, 59, 65, 75]. However, the underlying mechanisms are not yet completely understood. Smoking is thought to interfere with early healing events in the process of osseointegration and hence the consequences are usually recognized in the first year following implant placement. Lindquist *et al.* [59] reported that smokers demonstrated worse oral hygiene and displayed approximately 3 times greater bone loss after 10 years than non-smokers. Moreover, factors including heavy or unfavorable occlusal loading previously associated with increased peri-implant loss became more relevant with smoking [49].

Smoking and poor oral hygiene were found to be of greater influence on peri-implant bone loss than overload in long-term studies of patients treated with fixed partial dentures [24, 59]. Other studies showed that smoking and implant location in the maxilla were associated with an increased peri-implant marginal bone resorption [75].

It was however argued that smoking should not be an absolute contra-indication for implant therapy. Long-term heavy smokers would rather be informed of the increased risk of marginal bone loss at the implant site over the long-term, and eventually a higher risk of late implant failure [75]. In the 10-year follow-up study of Carlsson *et al.* [77], smoking was the most important factor affecting peri-implant bone loss in the mandible. A number of reviews of the literature emphasize smoking as a significant risk factor for a compromised prognosis of dental implants.

In agreement with that, the current study showed that smokers manifest a statistically significant higher degree of crestal bone loss when compared with non-smokers. Additionally, this effect was significantly enhanced when considered with other factors (TPS surface, anterior location and VBL < 1 mm). Interestingly, smoking even provoked other non-statistically significant factors to increase bone loss; these include opposing fixed or mixed occlusion, mandibular location, and standard implant collar heights (Tables 22, 23 and 25). These combinations were however still not statistically significant.

Despite that tobacco significantly influenced the implant success rates; it did not significantly lower the survival rate. Based on this result, one could argue that it cannot be considered a contra-indication for implant placement.

Oral hygiene and periodontal status effect

Oral hygiene and periodontal status are also of importance in regard to peri-implant bone loss and implant failure [55]. Isidor [39] found a progressive loss of radiographic bone and clinical probing depths at implants with enhanced plaque accumulation.

Although optimum dental hygiene was emphasized in the current study, patients were not enrolled in active and regular recall session. One might therefore assume that preliminary caries and pocket control – a procedure done systematically in the current study – is of a greater importance than strict post-treatment recall sessions. Consistency or frequency of recall and its effect on bone loss were not investigated in this study.

Implant location effect

Contradicting data in the literature have been reported on the effect of implant location (anterior versus posterior), on their success and survival rates. Weber *et al.* reported higher bone loss in the anterior region, although the size of the data was small [16]. Also, a 15-year prospective study demonstrated that implants placed in anterior segments showed higher bone loss than in the posterior segments [22]. Similarly, mesially placed implants showed more bone resorption than distally positioned implants, independently of implant surface roughness [14]. Lindquist *et al.* suggested that the more extensive bone loss around the anterior implants was a consequence of tensile forces, caused by loading of the posterior cantilever extensions and other biomechanical factors [24].

It is also debatable that implants placed in the posterior region show higher bone loss, considering that occlusal forces also increase because of the closeness of the temporomandibular joint. Therefore, all the posterior implants supporting partial prosthesis would experience more loading than those located in the anterior regions. Precisely, in a prospective 5-year study [20], the cumulative success rates for implants placed in molar sites were lower than mandibular and maxillary anterior regions. These differences reported for the anterior and posterior locations were attributed to bone quality and quantity (difficulty in achieving bicortical stabilization). Posterior regions were often characterized by unfavorable bone quality and reduced bone height, thus affecting bone loss and implant survival rate [78, 79]. Implants placed in the premolar or molar regions were generally shorter than those placed in the canine and incisor sites [64]. All of the above mentioned studies actually correspond to machined surfaced implants. With the introduction of roughened surfaces, research showed that posterior regions are no longer a risk factor.

Present results reported significant higher bone loss for implants placed in anterior regions. The combination of anterior region placement, tobacco use, and TPS-surfaced implant greatly increased bone loss. However, the results should be carefully interpreted as many of the anterior implants were usually placed deeper than those in the posterior region, to prevent exposure of the metal implant margin. Consequently, in addition to the crestal bone resorption which occurred for implants placed under standard conditions, bone adjacent to the polished implant surface was also lost. From a biological point of view, the subcrestal placement of the rough-smooth interface was consequently not recommended [27, 61].

Most of the posterior implants in the present study were short length. Due to the encouraging results in terms of CBL around posterior implants as opposed to anterior ones, short implant lengths could also be considered a valid treatment option and one with similar success rates to longer implants, especially in areas of reduced bone height.

Vestibular bone lamella width effect

The effect of VBL at implant placement on bone loss has not yet been reported in the literature. It is documented that a minimum width of 1 mm around implants, is required at placement to obtain optimum osseointegration and to prevent exposure of the implant threads following bone remodeling [5]. Primary stability was reported to be an even more significant factor for osseointegration [5, 56].

Bone that is less than 1 mm wide was thought to be more susceptible to resorption after surgery or following function. The rationale was that this minimal bone would fail to provide a sufficient matrix for the surrounding mesio-distal bone remodeling process, and would even enhance its resorption [12, 47, 77, 81]. Such findings should alarm clinicians while placing implants in anterior regions with a VBL less than 1 mm, since it may cause higher peri-implant bone loss and would therefore affect esthetic parameters, namely mesial and distal papillae.

There are reasons to suggest that over time this uneven outline of the marginal bone around dental implants was leveled out by a bone remodeling process and a reduction of the bone height at the proximal surfaces. Such an explanation is in agreement with findings reported by Carmagnola *et al.* [80] who, in a dog model, studied bone tissue reactions around implants placed in a compromised mandible. Following tooth extraction, the buccal bone plate was resected and a narrow ridge established. After 8 months of healing, implants were placed in the compromised site so that their lingual surfaces were invested in bone while about 4–5 mm of their buccal portion remained exposed. During the process of healing and during 4 months of function marked modeling and remodeling of the bone tissue around the implants took place. At the buccal surfaces some regrowth of bone occurred while at the lingual surfaces there was a substantial resorption of bone. As a result, the marginal level of osseointegration tended to become similar at all four aspects of the implants. This

finding could explain the greater proximal peri-implant bone loss around implants that have a buccal bone lamella that is smaller than 1 mm [51].

The width of the VBL showed a higher influence on CBL when combined with tobacco use, anterior arch location or TPS surfaced implants. The high value of CBL around implants having a VBL < 1 mm might be due to their anterior location which often involved a deeper fixture placement, this parameter was discussed previously. Moreover, implants with a VBL that is greater than 1 mm had more bone coronal to the R interface ("supra-boundary bone"), than those with a VBL that is less than 1mm.

Statistically non- significant factors affecting bone loss

All other investigated factors (implant diameter, opposing arch occlusion, maxillary versus mandibular jaw location, smooth collar height, fixed versus removable suprastructure, implant-implant or implant-tooth distance, and implant length) did not show a significant influence on bone remodeling in the current study. This could be attributed to the small number of patients who were concerned by these factors, errors in the radiographic analysis, or the actual negligible influence on bone remodeling and bone loss in the included patient sample. Nevertheless, when combined with other factors, some of these factors might influence bone levels. This will be discussed in the following paragraphs.

Implant diameter effect

Implant diameter is the distance between the peak of the widest thread and the same point on the opposite side of the implant. In contrast, implant diameter is distinct from the implant platform diameter, the latter being a measure of the interface of the implant connected with the abutment. Because a variety of implant widths and platforms are available, a wide-platform is not always related to an increased diameter of the implant thread. Implant diameter was not reported as a limiting factor in peri-implant bone loss [26, 82].

Reported advantages of using wide-diameter implants include: increased bone to implant contact, use as "rescue" implants in the case of site over-preparation during drilling, immediate placement in failure sites, reduction in abutment stresses and strain. The most obvious indication for wide-diameter implants (especially at the platform level) is for molar fixed rehabilitations [26].

Contrasting effects of implant diameter on success and survival rates were reported in the literature. Some studies report that 5 mm wide implants have higher failure rate than 3.75 or 4 mm wide implants because wider implants are often used in rescue procedures for failed implants [82]. Also, it has been reported that 5 mm wide implants developed for compromised situations had similar survival and success rates to standard-size Brånemark® implants.

Inversely, another study [83] showed that cumulative survival and success rates of small-diameter implants and standard-diameter implants were not statistically different ($p > 0.05$); although bone quality was a significant factor in failure, marginal bone loss was not influenced by the different implant diameters. The results suggested that small-diameter implants could be successfully used in the treatment of partially edentulous patients. Furthermore, no statistically significant relationship was observed between peri-implant bone loss and implant diameter [49, 84]. When considering wide neck ITI implants, a five-year life-table and radiographic analysis showed that these implants were highly predictable, with small prosthetic complications. The average bone loss measured at the two-year post-operative control was similar to standard implants [85].

No correlation between the different implant collar diameters on bone loss were noted in the present study. Wide neck implants did not show higher bone loss than smaller diameter implants. Moreover, combinations with other factors did not seem to have an effect.

Effect of opposing dentition

Occlusal overloading has been reported to be associated with increased bone loss and implant failure. This report was based on anecdotal observations supported by theoretical biomechanical theories, but was never proven in controlled studies in humans. Studies in monkeys demonstrated that overload could cause increase bone loss in some included implants [63]. Isidor [39] showed that overloaded implants had a decreased bone-to-implant contact. These overloaded implants also presented a smaller area in contact with mineralized bone tissue than non-loaded implants. Furthermore, once peri-implantitis has progressed, the control of occlusion and inflammation was probably not sufficient to promote the healing mechanism [18, 46, 65]. Implants with surrounding tissue inflammation probably deserve a greater care in avoiding overload. Again, these conclusions were drawn from studies done on machined surfaces.

Also, when reviewing the literature on the effect of the opposing dentition on bone loss, there was no conclusive answer to the question: did the prosthetic status in the opposing jaw influence the peri-implant bone loss and/or implant failure? Peri-implant bone loss may be enhanced in the jaw occluding with a fixed prosthesis in comparison with one occluding with a complete denture [86]. On the contrary, Carlsson *et al.* [77] did not experience such differences in peri-implant bone loss and suggested that the prosthesis in the opposing arch did not influence peri-implant bone loss.

The opposing dentition alone did not seem to influence bone loss in the present study. The combination of smoking and implants with a fixed or mixed opposing occlusion increased bone loss, unlike removable opposing dentition. This parameter deserves further analysis, since our results were not statistically significant.

Jaw location effect

Implant survival was lower in the maxilla than the mandible; this was attributed partially to a different bone quality in the maxilla [2, 14, 59, 65]. In the study of Carlsson *et al.* [77], all the failing – maxillary – implants (placed with 2-stage implant procedures) were lost during the healing period and not after the connection of the prosthesis. However, the same study reported similar peri-implant bone loss in both jaws.

Other studies revealed different results. A comparison between bone loss in the mandible and in the maxilla around 2-stage implants at abutment connection showed that a steady state was achieved after the first year of loading. The bone loss was 0.05 mm in the maxilla and 0.2 mm in the mandible for completely edentulous individuals wearing dentures [86]. Very few studies have been carried out on rough-surfaced implants [75], and in these no differences between maxilla and mandible were noted. A more recent study [87] showed that the implants located in the maxilla were associated with significantly higher bone loss.

No statistically significant differences in bone loss were noted around implants in the maxilla or in the mandible in the current study. However, the combination of smoking and mandibular location caused higher bone loss, unlike the combination of a maxillary location and a smoking subject. Further research on the influence of jaw location and bone conditions on oral implant outcomes are needed.

Effect of height of the smooth collar

It was shown that the height of the smooth implant collar has an effect on bone remodeling around the implant. Straumann® implants showed more marginal bone loss if the smooth part of the implant came into contact with the bone after a deeper placement [27]. This result led to the development of the “Esthetic” Plus line within the ITI Dental Implant System. The magnitude of initial bone remodeling around implants was dependant on the location of the rough-smooth border of the implant in an apico-coronal dimension [88]. The implant having the shortest smooth coronal collar showed no additional bone loss, while enabling deeper placement. Its use might reduce the risk of an exposed metal implant margin in areas of esthetic concern [81, 89].

The current study showed no significant difference between esthetic implants when compared with standard implants, in terms of crestal bone loss. This would be of interest as a deeper placement, especially in the anterior area, would jeopardize proximal bone.

Suprastructure effect

The systemic review of Berglundh *et al.* [12] has demonstrated that implants supporting overdentures exhibited higher frequencies of biological and technical complications

than implants with fixed reconstructions. A seven-year study reported similar survival rate for implants supporting single-tooth prostheses (95.6%), cantilever fixed partial prostheses (94.4%), fixed partial prostheses (96.1%), fixed complete prostheses (100%), and implant/tooth-supported prostheses (90.6%) and overdentures (95.7%) [89]. Mericske-Stern [62] observed that patterns of force transmission onto the implants were similar with a fixed complete denture and an overdenture connected to maxillary implants. The influence of mechanical and anatomical-prosthetic variables on peri-implant parameters was studied by several authors [48, 61–63, 89]. The type of the implant to denture attachments was shown to have little or no influence on the peri-implant parameters [90, 91]. The bar design did not significantly influence the occlusal force distribution pattern. Wyatt and Zarb [17] observed that implants supporting distal extensions prosthesis significantly increased bone loss in the first year of loading when compared to implants supporting prosthesis bounded by natural teeth.

Excessive marginal bone loss was explained by the overloading due to the lack of anterior contact and the presence of parafunctional activity [77]. It was shown that 70% of the occlusal forces were borne by the distal cantilevers and 30% by the implant-supported segment of the prosthesis on “Toronto bridges” or “Branemark bridges” [60, 61]. Biomechanical calculations and such results suggested that the most distal implants presented higher risk of bone loss because they were exposed to the largest forces, bending movements and stress concentrations. Subsequently, Nedir *et al.* did not experience lower survival rates of rough surfaced implants having a single unit distal extension [66].

In the present study, a removable suprastructure did not manifest greater peri-implant bone loss, despite the advanced age and presumed reduced dexterity of older patients. Overdentures do not represent a higher risk for the development of peri-implant lesions. Elderly patients with overdentures supported by attachments or bars can reasonably maintain healthy peri-implant conditions. The small sample of patients with a removable prosthesis might also explain the non-significance of the results.

Implant-tooth/implant-implant distance effect

Few investigations which assess the influence of the distance between implants or between implants and teeth in regard to bone loss are reported in the literature. Effects on the interdental papilla were thoroughly studied by Tarnow *et al.* [92]. This group observed that increased crestal bone loss would result in an increase in the distance between the base of the contact point of the adjacent crowns and the crest of bone. It is a proposed way to determine whether the papilla will be present or absent between two implants, and was previously reported between natural teeth. When multiple implants had to be placed in the esthetic zone, the use of small diameter implants might preserve at least 3 mm of bone at the implant-abutment level between them. Differences between

implant diameters did not however yield significant results when considering their effect on peri-implant bone remodeling, however, further research is needed.

No correlations between implant-implant or implant-tooth distances and mean bone level change were established in the present study.

Implant length effect

Reported studies on smooth-surfaced implants showed that short implants failed more frequently than longer ones [21, 60, 64, 95]. Historically, the use of short implant was not widely recommended because it was believed that occlusal forces might be dissipated over a large implant surface area to prevent excessive stresses at the interface [81]. However, finite element analysis (FEA) has shown that the occlusal forces are mainly distributed to the crestal bone, rather than evenly throughout the entire surface area of the implant interface [63]. Since masticatory forces were usually light and fleeting, they are normally well tolerated by the bone. This might explain why the implant length was not linearly related to biomechanical stability. Long term studies show a dramatic increase in failures for implants shorter than 7 mm in length, especially on machined surface implants, even more in type 4 bone [93].

Smoking, implant location and morphology, which were demonstrated to influence marginal bone loss, also associated with an increased failure rate with short implants [94]. Similarly, it was demonstrated that short implants, wide implants, implants supporting fixed prostheses, and implants placed in smokers were associated with a high CBL [49]. Implant length was the most significant factor in the maintenance of machined surfaced dental implants.

However, the introduction of rough surfaced implants has allowed a greater bone-to-implant contact; hence, higher success and survival rates were noted. Bernard *et al.* [94] suggested that the distinct magnitude of anchorage and the distinct loosening patterns registered for Brånemark® and Straumann® implant systems of different lengths might be related to the various surfaces. Greater torque forces were needed for rough implants of short length, unlike for implants with machined surfaces. Implants as short as 5 mm in length, with porous surface treatments, were introduced to replace possible sinus lift procedures [94]. Based on such observations of increased bone-to-implant contact on rough surfaces, private practitioners used short implants in various situations (e.g. in posterior maxilla with limited bone height or in posterior mandibular locations because of the proximity of the mandibular canal), they showed that short implants were as successful as long ones [64].

Renouard *et al.* [96] also demonstrated that the use of short implants could be considered for prosthetic rehabilitation of the severely resorbed maxilla as an alternative to more complicated surgical techniques; both implant failure rate and bone resorption over two years were not affected.

Mean marginal bone loss and gingival crevice probing depth associated with short or long implant lengths were

statistically comparable [50]. Accordingly, when considering the long-term multicenter evaluation of 2359 non-submerged Straumann® implants [21], the five year survival and success rates of 8 mm long implants did not differ significantly from the longer implants, despite the posterior placement of the shorter implants.

Peri-implant bone loss was quite similar for long and short implants in the present study, even when combined with other factors. Moreover, implants placed in the posterior area –that tended to be shorter for anatomical reasons- exhibited less bone loss than those placed in anterior areas. That confirmed the previous observation that length of rough-surfaced implant did not influence bone loss and implant success. This was a significant finding which might not only simplify surgical and planning procedures, but also might drastically expand the applications of implant therapy.

Other factors worthy of investigation

Further analysis on the effect of other factors on peri-implant bone remodeling might include prosthesis-related factors: the type of fixed suprastructure (single crown or fixed partial denture), nature of the fixed partial denture (two splinted crowns, a bridge, and a cantilever), misfit of the suprastructure, and crown to implant ratio [97].

The effect of the level of implant submersion and the delay of placement or loading might be also of interest. Patient periodontal status and number of visits to the hygienist are also shown to affect bone remodeling around implants. Other studies have observed that crestal bone level changes were correlated with the presence of a microgap even when a two-part implant (i.e. implant plus an abutment) was placed with a non-submerged technique [10]. When the microgap was located above the bone crest, less bone remodeling occurred; whereas when the microgap was placed below the bone crest, greater amounts of bone were lost. The lack of data in the studied population prevented the inclusion of the factors described above.

Conclusions

The conclusions of this study can be summarized in eight points. Some of these were already known and were thus confirmed; others were identified, in particular some interesting associations identifying some groups at risk:

- 1) The survival rate (99.2%) presented hereby compared well with related previously published studies.
- 2) Specific peri-implant bone loss beyond the smooth-rough implant interface was on average 1.2 mm, throughout a period of 5-6 years. Such value was in agreement with those reported in the literature on rough surface implants to date.

The radiographic method used for the evaluation could be described as unique: the bone level change was evaluated

from the smooth-rough implant interface considered as the baseline level; therefore, the measurements were done from 5-6 year post-operative radiographs only. The results confirm the reliability of the measurements.

- 3) The success rate (93%) was mainly based on the percentage of implants having CBL greater than 3 mm. These were considered as a higher failure risk group. Different criteria for success rate evaluations were used in different studies.
- 4) Two subpopulations within this study, presented higher failure risk groups, and should be monitored more closely and attentively when considering hygiene control and while establishing the treatment planning process:
 - 4.1 The first group represented 7% of the total population. It showed a CBL higher than 3 mm, which was considered as "alarming" by the study group.
 - 4.2 The second group included a population with the following factors: TPS-surfaced implants, anterior arch location, smokers, and VBL thinner than 1 mm at surgery. These factors – considered separately or combined – were associated with higher peri-implant CBL.
- 5) "Supra-boundary bone"; which is bone appearing above the rough-smooth interface was observed on 7.8% of the included implants. It was quite noticeable with SLA-surfaced implants, non-smoking subjects, implants located in a posterior arch, and implants with a VBL higher than 1 mm at surgery. These can be considered as "low failure risk groups".
- 6) Implants placed in the mandible, implants with an opposing mixed/fixed occlusion and implants supporting a removable suprastructure tended to cause higher bone loss, although results were not statistically significant.
- 7) Short implants showed a very limited bone loss, the difference in bone loss between short and long implants was not statistically significant. This confirmed the reliability of the use of short implants.
- 8) Systemic and continuous monitoring of peri-implant bone conditions along with the identification and control of associated risk factors are highly recommended for the diagnosis of peri-implant disease.

References

1. Lang NP, Wilson TG, Corbet EF. Biological complications with dental implants: their prevention, diagnosis and treatment. *Clin Oral Implants Res* 2000;11 Suppl 1:146-55.
2. Brånemark PI, Zarb GA, Albrektsson T. *Tissue-integrated prostheses* (pp 129-43). Quintessence, Chicago, 1985.
3. Adell R, Lekholm U, Rockler B, Brånemark PI. A 15-year study of osseointegrated implants in the treatment of the edentulous jaw. *Int J Oral Surg* 1981;10:387-416.
4. Adell R, Lekholm U, Rockler B, Brånemark PI, Lindhe J, Eriksson B, Sbordone L. Marginal tissue reactions at osseointegrated titanium fixtures (I). A 3-year longitudinal prospective study. *Int J Oral Maxillofacial Surg* 1986;15:39-52.
5. Albrektsson T, Zarb G, Worthington P, Eriksson AR. The long-term efficacy of currently used dental implants: a review and proposed criteria of success. *Int J Oral Maxillofac Implants* 1986;1:11-25.
6. De Smet E, Jacobs R, Gijbels F, Naert I. The accuracy and reliability of radiographic methods for the assessment of marginal bone level around oral implants. *Dentomaxillofac Radiol* 2002;31:176-81.
7. Buser D, Weber HP, Lang NP. Tissue integration of non-submerged implants. 1-year results of a prospective study with 100 ITI hollow-cylinder and hollow-screw implants. *Clin Oral Implants Res* 1990;1:33-40.
8. Hermann JS, Buser D, Schenk RK, Cochran DL. Crestal bone changes around titanium implants. A histometric evaluation of unloaded non-submerged and submerged implants in the canine mandible. *J Periodontol* 2000;71:1412-24.
9. Hermann JS, Schoolfield JD, Nummikoski PV, Buser D, Schenk RK, Cochran DL. Crestal bone changes around titanium implants: a methodological study comparing linear radiographic with histometric measurements. *Int J Oral Maxillofac Implants* 2001;16:475-85.
10. Hermann JS, Schoolfield JD, Schenk RK, Buser D, Cochran DL. Influence of the size of the microgap on crestal bone changes around titanium implants. A histometric evaluation of unloaded non-submerged implants in the canine mandible. *J Periodontol* 2001;72:1372-83.
11. Hermann JS, Cochran DL, Nummikoski PV, Buser D. Crestal bone changes around titanium implants. A radiographic evaluation of unloaded nonsubmerged and submerged implants in the canine mandible. *J Periodontol* 1997;68:1117-30.
12. Berglundh T, Persson L, Klinge B. A systematic review of the incidence of biological and technical complications in implant dentistry reported in prospective longitudinal studies of at least 5 years. *J Clin Periodontol* 2002;29 Suppl 3:197-212. 232-3.
13. Brägger U, Hugel-Pisoni C, Burgin W, Buser D. Correlations between radiographic, clinical and mobility parameters after loading of oral implants with fixed partial dentures. A 2-year longitudinal study. *Clin Oral Implants Res* 1996;7:230-9.
14. Zechner W, Trinkl N, Watzek G, Busenlechner D, Tepper G, Haas R, Watzek G. Radiologic follow-up of peri-implant bone loss around machine-surfaced and rough-surfaced interforaminal implants in the mandible functionally loaded for 3 to 7 years. *Int J Oral Maxillofac Implants* 2004;19:216-21.
15. Zybutz M, Rapoport D, Laurell L, Persson GR. Comparisons of clinical and radiographic measurements of inter-proximal vertical defects before and 1 year after surgical treatments. *J Clin Periodontol* 2000;27:179-86.
16. Weber HP, Crohin CC, Fiorellini JP. A 5-year prospective clinical and radiographic study of non-submerged dental implants. *Clin Oral Implants Res* 2000;11:144-53.
17. Wyatt CC, Zarb GA. Bone level changes proximal to oral implants supporting fixed partial prostheses. *Clin Oral Implants Res* 2002;13:162-8.
18. Fransson C, Lekholm U, Jemt T, Berglundh T. Prevalence of subjects with progressive bone loss at implants. *Clin Oral Implants Res* 2005;16:440-6.
19. Wenntröm J, Palmer R. Consensus report of session 3: clinical trails (pp 255-9). *Proceedings of the 3rd European workshop on Periodontology. Implant Dentistry*. Lang N, Karring T, Lindhe J eds. Quintessence, Berlin, 1999.

20. Jemt T, Lekholm U. Oral implant treatment in posterior partially edentulous jaws: a 5-year follow-up report. *Int J Oral Maxillofac Implants* 1993;8:635-40.
21. Buser D, Mericske-Stern R, Bernard JP, Behneke A, Behneke N, Hirt HP, Belsler UC, Lang NP. Long-term evaluation of non-submerged. ITI implants. Part 1: 8-year life table analysis of a prospective multi-center study with 2359 implants. *Clin Oral Implants Res* 1997;8:161-172.
22. Snauwaert K, Duyck J, van Steenberghe D, Quirynen M, Naert I. Time dependent failure rate and marginal bone loss of implant supported prostheses: a 15-year follow-up study. *Clin Oral Investig* 2000;4:13-20.
23. Mombelli A, Lang NP. The diagnosis and treatment of peri-implantitis. *Periodontol* 2000 1998;17:63-76.
24. Lindquist LW, Carlsson GE, Jemt T. A prospective 15-year follow-up study of mandibular fixed prostheses supported by osseointegrated implants. Clinical results and marginal bone loss. *Clin Oral Implants Res* 1996;7:329-36.
25. Grondahl K, Sundén S, Grondahl HG. Inter- and intraobserver variability in radiographic bone level assessment at Brånemark fixtures. *Clin Oral Implants Res* 1998;4:243-50.
26. Lee JH, Frias V, Lee KW, Wright RF. Effect of implant size and shape on implant success rates: a literature review. *Prosthet Dent* 2005;94:377-81.
27. Hämmerle CH, Brägger U, Burgin W, Lang NP. The effect of subcrestal placement of the polished surface of ITI implants on marginal soft and hard tissues. *Clin Oral Implants Res* 1996;7:111-9.
28. Hartmann GA, Cochran DL. Initial implant position determines the magnitude of crestal bone remodeling. *J Periodontol* 2004;75:572-7.
29. Bernard JP, Belsler UC, Martinet JP, Borgis SA. Osseointegration of Brånemark fixtures using a single-step operating technique. A preliminary prospective one-year study in the edentulous mandible. *Clin Oral Implants Res* 1995;6:122-9.
30. Pepelassi EA, Diamanti-Kipiotti A. Selection of the most accurate method of conventional radiography for the assessment of periodontal osseous destruction. *J Clin Periodontol* 2000;27:179-86.
31. Romeo E, Lops D, Amorfini L, Chiapasco M, Ghisolfi M, Vogel G. Clinical and radiographic evaluation of small-diameter (3.3-mm) implants followed for 1-7 years: a longitudinal study. *Clin Oral Implants Res* 2006;17:139-48.
32. Rosling B, Hollender L, Nyman S, Olsson G. A radiographic method for assessing changes in alveolar bone height following periodontal therapy. *J Clin Periodontol* 1975;2:211-7.
33. Zechner W, Watzak G, Gahleitner A, Busenlechner D, Tepper G, Watzek G. Rotational panoramic versus intraoral rectangular radiographs for evaluation of peri-implant bone loss in the anterior atrophic mandible. *Int J Oral Maxillofac Implants* 2003;18:873-8.
34. Sewerin IP. Errors in radiographic assessment of marginal bone height around osseointegrated implants. *Scand J Dent Res* 1990;98:428-33.
35. Benn DK. Limitations of the digital image subtraction technique in assessing alveolar bone crest changes due to misalignment errors during image capture. *Dentomaxillofac Radiol* 1990;19:97-104.
36. Wahl G, Lang H. Deformation at the implant interface to prosthetic superstructure: an interferometric approach. *Clin Oral Implants Res* 2004;15:233-8.
37. Spiekermann H, Jansen VK, Richter EJ. A 10-year follow-up study of IMZ and TPS implants in the edentulous mandible using bar-retained overdentures. *Int J Oral Maxillofac Implants* 1995;10:231-43.
38. Quirynen M, van Steenberghe D, Jacobs R, Schotte A, Darius P. The reliability of pocket probing around screw-type implants. *Clin Oral Implants Res* 1991;2:186-92.
39. Isidor F. Clinical probing and radiographic assessment in relation to the histologic bone level at oral implants in monkeys. *Clin Oral Implants Res* 1997;8:255-64.
40. Ericsson I, Lindhe J. Probing depth at implants and teeth. An experimental study in the dog. *J Clin Periodontol* 1993;20:623-7.
41. Akesson L, Hakansson J, Rohlin M. Comparison of panoramic and intraoral radiography and pocket probing for the measurement of the marginal bone level. *J Clin Periodontol* 1992;19:326-32.
42. Lofthag-Hansen S, Lindh C, Petersson A. Radiographic assessment of the marginal bone level after implant treatment: a comparison of periapical and Scanora detailed narrow beam radiography. *Dentomaxillofac Radiol* 2003;32:97-103.
43. Battenberg RH, Meijer HJ, Raghoobar GM, Van Port RP, Boering G. Mandibular overdentures supported by two Brånemark, IMZ or ITI implants. A prospective comparative preliminary study: one-year results. *Clin Oral Implants Res* 1998;9:374-383.
44. Fiorellini JP, Buser D, Paquette DW, Williams RC, Haghghi D, Weber HP. A radiographic evaluation of bone healing around submerged and non-submerged dental implants in beagle dogs. *J Periodontol* 1999;70:248-54.
45. Heydenrijk K, Raghoobar GM, Meijer HJ, Stegenga B. Clinical and radiologic evaluation of 2-stage IMZ implants placed in a single-stage procedure: 2-year results of a prospective comparative study. *Int J Oral Maxillofac Implants* 2003;18:424-32.
46. Karoussis K, Muller S, Salvi G, Heitz-Mayfield L, Brägger U, Lang N. Association between periodontal and peri-implant conditions ; a 10 year prospective study. *Clin Oral Impl Res* 2004;15:1-7.
47. Meijer HJ, Heijdenrijk K, Raghoobar GM. Mucosal and radiographic aspects during the healing period of implants placed in a one-stage procedure. *Int J Prosthodont* 2003;16:397-402.
48. Mericske-Stern R, Oetterli M, Kiener P, Mericske E. A follow-up study of maxillary implants supporting overdentures. clinical and radiographic results. *Int J Oral Maxillofac Implants* 2002;17:678-86.
49. Penarrocha M, Palomar M, Sanchis JM, Guarinos J, Balaguer J. Radiologic study of marginal bone loss around 108 dental implants and its relationship to smoking, implant location, and morphology. *Int J Oral Maxillofac Implants* 2004;19:861-78.
50. Romeo E, Ghisolfi M, Rozza R, Chiapasco M, Lops D. Short (8mm) dental implants in the rehabilitation of partial and complete edentulism: a 3 to 14 year longitudinal study. *Int J Prosthodont* 2006;19:586-92.
51. Blanes RJ, Bernard JP, Blanes ZM, Belsler UC. A 10-year prospective study of ITI dental implants placed in the posterior region: I Clinical and radiographic results. *Clin Oral Implants Res* 2007;18:699-706.
52. Von Wowern N. In vivo measurement of bone mineral content of mandibles by dual photon absorptiometry. *Scand J Dent Res* 1985;93:162-8.
53. Brägger U, Pasquali L, Kornman KS. Remodeling of interdental alveolar bone after periodontal flap procedures assessed

- by means of computer-assisted densitometric image analysis (CADIA). *J Clin Periodontol* 1988;15:558-4.
54. Cochran DL, Hermann JS, Schenk RK, Higginbottom FL, Buser D. Biologic width around titanium implants. A histometric analysis of the implant-to-gingival junction around unloaded and loaded nonsubmerged implants in the canine mandible. *J Periodontol* 1997;68:186-98.
 55. Smith DE, Zarb GA. Criteria for success of osseointegrated endosseous implants. *J Prosthet Dent* 1989;62:567-72.
 56. Abrahamsson I, Zitzmann NU, Berglundh T, Wennerberg A, Lindhe J. Bone and soft tissue integration to titanium implants with different surface topography: an experimental study in the dog. *Int J Oral Maxillofac Implants* 2001;16:323-33.
 57. Meriscke-Stern R, Venetz E, Fahrlander F, Burgin W. In vivo force measurements on maxillary implants supporting a fixed prosthesis or an overdenture: a pilot study. *J Prosthet Dent* 2000;84:535-47.
 58. Keller W, Brägger U, Mombelli A. Peri-implant microflora of implants with cemented and screw retained suprastructures. *Clin Oral Implants Res* 1998;9:209-17.
 59. Lindquist LW, Carlsson GE, Jemt T. Association between marginal bone loss around osseointegrated mandibular implants and smoking habits: a 10-year follow-up study. *J Dent Res* 1997;76:1667-74.
 60. Lum LB. A biomechanical rationale for the use of short implants. *J Oral Implantol* 1991;17:126-31.
 61. Mailath G, Stoiber B, Watzek G, Matejka M. Bone resorption at the entry of osseointegrated implants - a biomechanical phenomenon. Finite element study. *Z Stomatol* 1989;86:207-16.
 62. Meriscke-Stern R. Force distribution on implants supporting overdentures: the effect of distal bar extensions. A 3-D in vivo study. *Clin Oral Implants Res* 1997;8:142-51.
 63. Miyata T, Kobayashi Y, Araki H, Ohto T, Shin K. The influence of controlled occlusal overload on peri-implant tissue. Part 4: a histologic study in monkeys. *Int J Oral Maxillofac Implants* 2002;17:384-90.
 64. Nedir R, Bischof M, Briaux JM, Beyer S, Szmukler-Moncler S, Bernard JP. A 7-year life table analysis from a prospective study on ITI implants with special emphasis on the use of short implants. Results from a private practice. *Clin Oral Implants Res* 2004;15:150-7.
 65. Esposito M, Hirsch JM, Lekholm U, Thomsen P. Biological factors contributing to failures of osseointegrated oral implants. (I) Success criteria and epidemiology. *Eur J Oral Sci* 1998;106:527-51.
 66. Nedir R, Bischof M, Szmukler-Moncler S, Belser UC, Samson J. Prosthetic complications with dental implants: from an up-to-8-year experience in private practice. *Int J Oral Maxillofac Implants* 2006;21:919-28.
 67. Scharf DR, Tarnow DP. Success rates of osseointegration for implants placed under sterile versus clean conditions. *J Periodontol* 1993;64:954-6.
 68. Buser D, von Arx T. Surgical procedures in partially edentulous patients with ITI implants. *Clin Oral Implants Res* 2000;11 Suppl 1:83-100.
 69. Hämmerle CH, Ingold HP, Lang NP. Evaluation of clinical and radiographic scoring methods before and after initial periodontal therapy. *J Clin Periodontol* 1990;17:255-63.
 70. Arlin ML. Survival and success of sandblasted, large-grit, acid-etched and titanium plasma-sprayed implants: a retrospective study. *J Can Dent Assoc* 2007;73:821.
 71. Cochran DL, Buser D, ten Bruggenkate CM, Weingart D, Taylor TM, Bernard JP, Peters F, Simpson JP. The use of reduced healing times on ITI implants with a sandblasted and acid-etched (SLA) surface: early results from clinical trials on ITI SLA implants. *Clin Oral Implants Res* 2002;13:144-53.
 72. Cochran DL, Schenk RK, Lussi A, Higginbottom FL, Buser D. Bone response to unloaded and loaded titanium implants with a sandblasted and acid-etched surface: a histometric study in the canine mandible. *J Biomed Mater Res* 1998;40:1-11.
 73. Bornstein MM, Schmid B, Belser UC, Lussi A, Buser D. Early loading of non-submerged titanium implants with a sandblasted and acid-etched surface. 5-year results of a prospective study in partially edentulous patients. *Clin Oral Implants Res* 2005;16:631-8.
 74. Rocuzzo M, Wilson TG. A prospective study evaluating a protocol for 6 weeks' loading of SLA implants in the posterior maxilla. *Clin Oral Implants Res* 2002;13:502-7.
 75. DeLuca S, Zarb G. The effect of smoking on osseointegrated dental implants. Part II: Peri-implant bone loss. *Int J Prosthodont* 2006;19:560-6.
 76. Gruica B, Wang HY, Lang NP, Buser D. Impact of IL-1 genotype and smoking status on the prognosis of osseointegrated implants. *Clin Oral Implants Res* 2004;15:393-400.
 77. Carlsson GE, Lindquist LW, Jemt T. Long-term marginal peri-implant bone loss in edentulous patients. *Int J Prosthodont* 2000;13:295-302.
 78. Ozkan Y, Ozcan M, Akoglu B, Ucankale M, Kulak-Ozkan Y. Three-year treatment outcomes with three brands of implants placed in the posterior maxilla and mandible of partially edentulous patients. *J Prosthet Dent* 2007;97:78-84.
 79. Quirynen M, Naert I, van Steenberghe D, Nys L. A study of 589 consecutive implants supporting complete fixed prostheses. Part I: Periodontal aspects. *J Prosthet Dent* 1992;68:655-63.
 80. Carmagnola D, Araujo M, Berglundh T, Albrektsson T, Lindhe J. Bone tissue reaction around implants placed in a compromised jaw. *J Clin Periodontol* 1999;26:629-35.
 81. Brägger U, Hafeli U, Huber B, Hammerle CH, Lang NP. Evaluation of postsurgical crestal bone levels adjacent to non-submerged dental implants. *Clin Oral Implants Res* 1998;9:218-24.
 82. Renouard F, Arnoux JP, Sarment DP. Five-mm-diameter implants without a smooth surface collar: report on 98 consecutive placements. *Int J Oral Maxillofac Implants* 1999;14:101-7.
 83. Ivanoff CJ, Grondahl K, Sennerby L, Bergstrom C, Lekholm U. Influence of variations in implant diameters: a 3- to 5-year retrospective clinical report. *Int J Oral Maxillofac Implants* 1999;14:173-80.
 84. Block MS, Delgado A, Fontenot MG. The effect of diameter and length of hydroxylapatite-coated dental implants on ultimate pullout force in dog alveolar bone. *J Oral Maxillofac Surg* 1990;48:174-8.
 85. Bischof M, Nedir R, Abi Najm S, Szmukler-Moncler S, Samson J. A five-year life-table analysis on wide neck ITI implants with prosthetic evaluation and radiographic analysis: results from a private practice. *Clin Oral Implants Res* 2006;17:512-20.
 86. Tallgren A. The continuing reduction of the residual alveolar ridges in complete denture wearers: a mixed-longitudinal study covering 25 years. *J Prosthet Dent* 1972;27:120-32.
 87. Penarrocha M, Guarinos J, Sanchis JM, Balaguer J. A retrospective study (1994-1999) of 441 ITI(r) implants in 114 patients followed-up during an average of 2.3 years. *Med Oral* 2002;7:144-55.

88. Hanggi MP, Hanggi DC, Schoolfield JD, Meyer J, Cochran DL, Hermann JS. Crestal bone changes around titanium implants. Part I: A retrospective radiographic evaluation in humans comparing two non-submerged implant designs with different machined collar lengths. *J Periodontol* 2005;76:791-802.
89. Romeo E, Lops D, Margutti E, Ghisolfi M, Chiapasco M, Vogel G. Long-term survival and success of oral implants in the treatment of full and partial arches: a 7-year prospective study with the ITI dental implant system. *Int J Oral Maxillofac Implants* 2004;19:247-59.
90. Kiener P, Oetterli M, Mericske E, Mericske-Stern R. Effectiveness of maxillary overdentures supported by implants: maintenance and prosthetic complications. *Int J Prosthodont* 2001;14:133-40.
91. Oetterli M, Kiener P, Mericske-Stern R. A longitudinal study on mandibular implants supporting an overdenture: the influence of retention mechanism and anatomic-prosthetic variables on peri-implant parameters. *Int J Prosthodont* 2001;14:536-42.
92. Tarnow DP, Cho SC, Wallace SS. The effect of inter-implant distance on the height of inter-implant bone crest. *J Periodontol* 2000;71:546-9.
93. Ivanoff CJ, Grondahl K, Bergstrom C, Lekholm U, Brånemark PI. Influence of bicortical or monocortical anchorage on maxillary implant stability: a 15-year retrospective study of Brånemark system implants. *Int J Oral Maxillofac Implants* 2000;15:103-10.
94. Bernard JP, Szmukler-Moncler S, Pessotto S, Vazquez L, Belser UC. The anchorage of Brånemark and ITI implants of different lengths. I. An experimental study in the canine mandible. *Clin Oral Implants Res* 2003;14:593-600.
95. Deporter D, Todescan R, Caudry S. Simplifying management of the posterior maxilla using short, porous-surfaced dental implants and simultaneous indirect sinus elevation. *Int J Periodont Restorat Dent* 2000;20:476-85.
96. Renouard F, Nisand D. Short implants in the severely resorbed maxilla: a 2-year retrospective clinical study. *Clin Implant Dent Relat Res* 2005;7 Suppl 1:104-10.
97. Rokni S, Todescan R, Watson P, Pharoah M, Adegbembo AO, Deporter D. An assessment of crown-to-root ratios with short sintered porous-surfaced implants supporting prostheses in partially edentulous patients. *Int J Oral Maxillofac Implants* 2005;20:69-76.